

Bosonic realization of toroidal Lie algebras of classical types

Naihuan Jing*, Kailash C. Misra, and Chongbin Xu

ABSTRACT. Generalizing Feingold-Frenkel's construction we use Weyl bosonic fields to construct toroidal Lie algebras of types A_n, B_n, C_n and D_n of level $-1, -2, -1/2$ and -2 respectively. In particular, our construction also gives new bosonic construction for the orthogonal Lie algebras in the cases of affine Lie algebras.

1. Introduction

Toroidal Lie algebras are natural generalization of the affine Kac-Moody algebras [MRY] that enjoy many similar interesting features. Let $\mathfrak{g}$ be a finite-dimensional complex simple Lie algebra of type X_n , and $R = \mathbb{C}[s, s^{-1}, t, t^{-1}]$ be the ring of Laurent polynomials in commuting variables s and t . By definition a 2-toroidal Lie algebra of type X_n is a perfect central extension of the iterated loop algebra $\mathfrak{g} \otimes R$, and can be realized as certain homomorphic image of the universal central extension $T(X_n) = (g \otimes R) \oplus \Omega_R/dR$, where Ω_R/dR is Kähler differentials of R modulo the exact forms. The center contains two special elements c_0, c_1 . A module of $T(X_n)$ is called a level- (k_0, k_1) module if the standard pair of central elements (c_0, c_1) acts as (k_0, k_1) for some complex numbers k_0 and k_1 . In this work we will focus on modules with $k_0 \neq 0$.

Two-toroidal Lie algebras resemble affine Lie algebras in many aspects. Their main source of representations are vertex operators (cf. [MRY, B, BBS]). In [T] the toroidal Lie algebra of type B_n was constructed by fermionic operators (see also [JMg]), and in [G] two constructions were given for the extended affine Lie algebras of type A_n . In [FJW] the first author and collaborators used McKay correspondence to realize level one representation of toroidal Lie algebras of simply laced types. In the recent paper [JM] a unified fermionic construction of toroidal Lie algebras of classical types was given by extending Feingold-Frenkel realization coupled with new ghost fields. We pointed out there that the fermionic construction can not be directly generalized to symplectic toroidal Lie algebras as one needs bosons to realize the long root vector $2\varepsilon_n$.

2000 *Mathematics Subject Classification.* Primary: 17B60, 17B67, 17B69. Secondary: 17A45, 81R10.

Key words and phrases. Toroidal algebras, Weyl algebras, vertex operators, representations.

* Corresponding author.

Jing acknowledges the support of NSA grant H98230-06-1-0083 and NSFC grant 10728102, and Misra acknowledges the support of NSA grant H98230-08-0080.

In this paper we give a unified Weyl bosonic construction of all classical toroidal Lie algebras. Our work in types C and A are toroidal analog of Feingold-Frenkel construction [FF] for the affine Lie algebras (see also [L] for a recent treatment of Feingold-Frenkel construction). We also construct bosonic realizations for types D and B . This includes, as special cases, new constructions (at level -2) for orthogonal affine Lie algebras. The main idea is again similar to that of [JM] to construct certain operators corresponding to the special nodes in the affine Dynkin diagrams. This novelty is special for the cases of toroidal Lie algebras. In a sense we have also obtained field operators corresponding to imaginary root vectors.

The Weyl bosonic construction for orthogonal toroidal Lie algebras is not directly obtained by mimicking that of type A or C and we have to use new embedding of the Lie algebras into type A to achieve the goal, this partly explains why this construction has been missed in previous available constructions of affine Lie algebras. Like the Neveu-Schwarz and Ramond fields in the fermionic case, our field operators have two forms indexed by half-integers and integers respectively.

The structure of the paper is as follows. In section 2 we define the toroidal Lie algebra, and state MRY-presentation [MRY] of the toroidal algebra in terms of generators and relations. In section 3 we start with a finite rank lattice with an anti-symmetric bilinear form and define a Fock space and some bosonic field (vertex) operators, which in turn give level -1 representations of the toroidal Lie algebras of type A_n , level $-1/2$ modules for type C , and level -2 modules for types B_n, D_n . The proof is an extensive analysis of the operator product expansions for the field operators. We also include the verification of the Serre relations.

2. Toroidal Lie Algebras

A special quotient algebra of the toroidal Lie algebra $T(X_n)$ is the double affine algebra, denoted by $T_0(X_n)$, that is the toroidal Lie algebra of type X_n with a two dimensional center. The double affine algebra is the quotient of $T(X_n)$ modulo all the central elements with degree other than zero. In fact, $T_0(X_n)$ has the following realization

$$T_0(X_n) = (\mathfrak{g} \otimes R) \oplus \mathbb{C}c_0 \oplus \mathbb{C}c_1,$$

where R is a ring and the Lie product is

$$[x \otimes g_1, y \otimes g_2] = [x, y] \otimes g_1 g_2 + \Phi(g_2 \partial_s g_1) c_0 + \Phi(g_2 \partial_t g_1) c_1$$

for all $x, y \in \mathfrak{g}$, $g_1, g_2 \in R$, where Φ is a linear functional on R defined by $\Phi(s^k t^m) = 0$, if $(k, m) \neq (0, 0)$ and $\Phi(s^k t^m) = 1$, if $(k, m) = (0, 0)$ for all $k, m \in \mathbb{Z}$.

For our purpose we will need the formal power series in variables z, w . In particular we will need the formal delta functions $\delta(z - w) = \sum_{n \in \mathbb{Z}} z^{-n-1} w^n$. This could be understood as follows.

$$\begin{aligned} \delta(z - w) &= \iota_{z,w}((z - w)^{-1}) + \iota_{w,z}((w - z)^{-1}), \\ \partial_w \delta(z - w) &= \iota_{z,w}((z - w)^{-2}) - \iota_{w,z}((w - z)^{-2}), \end{aligned}$$

where $\iota_{z,w}$ means expansion when $|z| > |w|$. For simplicity in the following we will drop $\iota_{z,w}$ if it is clear from the context.

For $n \geq 1$ let $(a_{ij})_{n+1 \times n+1}$ be the generalized Cartan matrix of the affine algebra $X_n^{(1)}$, and $Q := \mathbb{Z}\alpha_0 \oplus \mathbb{Z}\alpha_1 \oplus \cdots \oplus \mathbb{Z}\alpha_n$ its root lattice. The toroidal Lie algebra $T(X_n)$ [MRY] is the Lie algebra generated by ℓ , $\alpha_i(k)$, and $x_k(\pm\alpha_i)$ for $i = 0, 1, \dots, n$, $k \in \mathbb{Z}$ with the following relations:

- (R0) $[\not\partial, \alpha_i(z)] = 0 = [\not\partial, x(\pm\alpha_i, z)];$
(R1) $[\alpha_i(z), \alpha_j(w)] = (\alpha_i|\alpha_j)\partial_w\delta(z-w)\not\partial;$
(R2) $[\alpha_i(z), x(\pm\alpha_j, w)] = \pm(\alpha_i|\alpha_j)x(\pm\alpha_j, w)\delta(z-w);$
(R3) $[x(\alpha_i, z), x(-\alpha_j, w)] = \delta_{ij}\frac{2}{(\alpha_i|\alpha_j)}\{\alpha_i(w)\delta(z-w) + \partial_w\delta(z-w)\not\partial\};$
(R4) $[x(\alpha_i, z), x(\alpha_i, w)] = 0 = [x(-\alpha_i, z), x(-\alpha_i, w)];$
(S1) $\text{adx}(\pm\alpha_i, z_1)x(\pm\alpha_j, z_2) = 0, \quad \text{for } a_{ij} = 0$
(S2) $(\text{adx}(\pm\alpha_i, z_1))(\text{adx}(\pm\alpha_i, z_2))x(\pm\alpha_j, z_3) = 0, \text{ if } a_{ij} = -1$
(S3) $(\text{adx}(\pm\alpha_i, z_1))(\text{adx}(\pm\alpha_i, z_2))(\text{adx}(\pm\alpha_i, z_3))x(\pm\alpha_j, z_4) = 0, \text{ if } a_{ij} = -2.$
- where we have used the follow power series:

$$\alpha_i(z) = \sum_{n \in \mathbb{Z}} \alpha_i(n) z^{-n-1},$$

$$x(\pm\alpha_i, z) = \sum_{n \in \mathbb{Z}} x_n(\pm\alpha_i) z^{-n-1}.$$

We remark that the 2-toroidal Lie algebra of type X_n includes two affine Lie algebras of type $X_n^{(1)}$ as subalgebras.

3. Representations of the Toroidal Algebra

In this section we give two unified bosonic realizations for the toroidal Lie algebra of classical types A_{n-1}, B_n, C_n and D_n using bosonic analogs of Neveu-Schwarz and Ramond fermionic fields.

Let ε_i ($i = 1, \dots, n+1$) be a set of orthonormal basis of the vector space $\mathbb{C}^{n+1}$ equipped with the inner product $(\cdot | \cdot)$ such that

$$(3.1) \quad (\varepsilon_i | \varepsilon_j) = \delta_{ij},$$

Let $P_0 = \mathbb{Z}\varepsilon_1 \oplus \dots \oplus \mathbb{Z}\varepsilon_n$ be a sublattice of rank n , and let ε'_0 be an external unit vector orthogonal to ε_i and introduce $\bar{c} = \frac{1}{\sqrt{2}}(\varepsilon'_0 + i\varepsilon_{n+1})$ correspond to the null vector δ and $\bar{d} = \frac{1}{\sqrt{2}}(\varepsilon'_0 - i\varepsilon_{n+1})$ be the dual gradation operator. Then

$$(3.2) \quad (\bar{c}|\bar{c}) = (\bar{c}|\varepsilon_i) = 0$$

$$(3.3) \quad (\bar{d}|\bar{d}) = (\bar{d}|\varepsilon_i) = 0, \quad (\bar{c}|\bar{d}) = 1$$

for $i = 1, \dots, n$.

The simple roots for the classical finite dimensional Lie algebras can be realized simply by defining the simple roots as follows:

$$\begin{aligned} \alpha_1 &= \varepsilon_1 - \varepsilon_2, \dots, \alpha_{n-1} = \varepsilon_{n-1} - \varepsilon_n; \text{ for } A_{n-1}. \\ \alpha_1 &= \varepsilon_1 - \varepsilon_2, \dots, \alpha_{n-1} = \varepsilon_{n-1} - \varepsilon_n, \alpha_n = \varepsilon_n; \text{ for } B_n. \\ \alpha_1 &= \frac{1}{\sqrt{2}}(\varepsilon_1 - \varepsilon_2), \dots, \alpha_{n-1} = \frac{1}{\sqrt{2}}(\varepsilon_{n-1} - \varepsilon_n), \alpha_n = \sqrt{2}\varepsilon_n; \text{ for } C_n. \\ \alpha_1 &= \varepsilon_1 - \varepsilon_2, \dots, \alpha_{n-1} = \varepsilon_{n-1} - \varepsilon_n, \alpha_n = \varepsilon_{n-1} + \varepsilon_n; \text{ for } D_n. \end{aligned}$$

Then the set of positive roots are:

$$\Delta_+ = \begin{cases} \{\varepsilon_i - \varepsilon_j | 1 \leq i < j \leq n\}, & \text{Type } A_{n-1} \\ \{\varepsilon_i, \varepsilon_i \pm \varepsilon_j | 1 \leq i < j \leq n\}, & \text{Type } B_n \\ \{\sqrt{2}\varepsilon_i, \frac{1}{\sqrt{2}}(\varepsilon_i \pm \varepsilon_j) | 1 \leq i < j \leq n\}, & \text{Type } C_n \\ \{\varepsilon_i \pm \varepsilon_j | 1 \leq i < j \leq n\}, & \text{Type } D_n. \end{cases}$$

The highest (long) root α_{max} for each type is given as follows:

$$\alpha_{max} = \begin{cases} \varepsilon_1 - \varepsilon_n, & \text{Type } A_{n-1} \\ \varepsilon_1 + \varepsilon_2, & \text{Type } B_n \text{ or } D_n. \\ \sqrt{2}\varepsilon_1 & \text{Type } C_n \end{cases}$$

We further introduce the element

$$\alpha_0 = \bar{c} - \alpha_{max}$$

in the lattice and then define $\beta = -\bar{c} + \varepsilon_1$ for type ABD , and $\beta = -\sqrt{2}\bar{c} + \varepsilon_1$ for type C . Then we have

$$\alpha_0 = \varepsilon_n - \beta \text{ for } A_{n-1}; \quad -\beta - \varepsilon_2 \text{ for } B_n, D_n; \quad \text{or } -\frac{1}{\sqrt{2}}(\beta + \varepsilon_1) \text{ for } C_n.$$

Note that $(\beta|\beta) = 1, (\beta|\varepsilon_i) = \delta_{1i}$.

Then $P = \mathbb{Z}\bar{c} \oplus \mathbb{Z}\varepsilon_1 \oplus \cdots \oplus \mathbb{Z}\varepsilon_n$ and $Q = \mathbb{Z}\bar{c} \oplus \mathbb{Z}\alpha_1 \oplus \cdots \oplus \mathbb{Z}\alpha_k = \mathbb{Z}\alpha_0 \oplus \cdots \oplus \mathbb{Z}\alpha_k$ ($k = n-1$ for type A_{n-1} and $k = n$ for types B_n, C_n, D_n) are the weight lattice and root lattice for the corresponding affine Lie algebra, and $P_0 = \mathbb{Z}\varepsilon_1 \oplus \cdots \oplus \mathbb{Z}\varepsilon_n$ and $Q_0 = \mathbb{Z}\alpha_1 \oplus \cdots \oplus \mathbb{Z}\alpha_k$ ($k = n-1$ for type A_{n-1} and $k = n$ for types B_n, C_n, D_n) are the weight lattice and root lattice for the simple Lie algebras A_{n-1}, B_n, C_n, D_n . Then $(\alpha_i|\alpha_j) = d_i a_{ij}$, where a_{ij} are the entries of the affine Cartan matrix of type $(ABCD)^{(1)}$, and the d_i 's are given by:

$$(d_0, d_1, \dots, d_k) = \begin{cases} \{1, 1, \dots, 1, 1\}, & k = n-1, \text{Type } A_{n-1} \\ \{1, 1, \dots, 1, \frac{1}{2}\}, & k = n, \text{Type } B_n \\ \{1, \frac{1}{2}, \dots, \frac{1}{2}, 1\}, & k = n, \text{Type } C_n \\ \{1, 1, \dots, 1, 1\}, & k = n, \text{Type } D_n. \end{cases}$$

Note that the bilinear form $(\ | \)$ is non-degenerate on the affine root lattice or the span of $\{\beta\} \cup \{\alpha_i\}$ ($i \geq 1$).

We introduce infinite dimensional Weyl algebras as follows. Let $\mathbf{Z} = \mathbb{Z}$ or $\mathbf{Z} = \mathbb{Z} + 1/2$. Let $P_{\mathbb{C}} = P \otimes \mathbb{C}$ be the $\mathbb{C}$ -vector space spanned by $\bar{c}$ and $\varepsilon_i, 1 \leq i \leq n$ in various types. To combine all types we will also introduce another set of orthonormal vectors $\varepsilon_{\bar{i}}, 1 \leq i \leq n+1$ such that

$$(3.4) \quad (\varepsilon_{\bar{i}}|\varepsilon_{\bar{j}}) = \delta_{ij},$$

and we denote by $\bar{P}_{\mathbb{C}}$ the $\mathbb{C}$ -space spanned by $\varepsilon_{\bar{i}}$. We also define $\bar{\beta} = -\bar{c} + \varepsilon_{\bar{1}}$, and $\alpha_{\bar{0}} = \bar{c} - \alpha_{\overline{max}}$, where $\alpha_{\overline{max}} = \varepsilon_{\bar{1}} + \varepsilon_{\bar{2}}$. The special vectors are only needed in types B_n, D_n and $\varepsilon_{\overline{n+1}}$ is only needed for type B_n . We define $\mathcal{C} = \mathcal{C}_1 \oplus \mathcal{C}_2$, where both subspaces $\mathcal{C}_1 = P_{\mathbb{C}} \oplus \bar{P}_{\mathbb{C}}$ and $\mathcal{C}_2 = P_{\mathbb{C}}^* \oplus \bar{P}_{\mathbb{C}}^*$ are maximal isotropic subspaces, where we define $\langle P, \bar{P} \rangle = 0$. We define the natural anti-symmetric bilinear form on $\mathcal{C}$ by

$$(3.5) \quad \langle b^*, a \rangle = -\langle a, b^* \rangle = (a|b), \quad \langle a, b \rangle = \langle a^*, b^* \rangle = 0, \quad a, b \in \mathcal{C}_1$$

The Weyl algebra $W(P)$ is the unital algebra generated by the elements $a(k)$ and $a^*(k)$, where $a \in \mathcal{C}_1, a^* \in \mathcal{C}_2$, and $k \in \mathbf{Z}$ subject to the relations:

$$[u(k), v(l)] = \langle u, v \rangle \delta_{k, -l}, \quad u, v \in \mathcal{C}.$$

The representation space is the infinite dimensional vector space

$$V = \bigotimes_{a_i} \left(\bigotimes_{k \in \mathbf{Z}_+} \mathbb{C}[a_i(-k)] \bigotimes_{k \in \mathbf{Z}_+} \mathbb{C}[a_i^*(-k)] \right)$$

where a_i runs through any basis in P and $\overline{P}$, say $\bar{c}_i$, ε_i 's and ε_i^- 's.

The Weyl algebra acts on the space V by the usual action: $a(-k)$ acts as an creation operator and $a(k)$ as an annihilation operator. We remark that using $-\langle \cdot, \cdot \rangle$ as the form on $\mathcal{C}$ is equivalent to switching $\mathcal{C}_1$ and $\mathcal{C}_2$. As the symplectic form $\langle \cdot, \cdot \rangle$ is non-degenerate, the Fock space V is an irreducible module for the Weyl algebra.

For any bosonic fields

$$u(z) = \sum_{n \in \mathbf{Z}} u(n) z^{-n-1/2}, \quad v(z) = \sum_{n \in \mathbf{Z}} v(n) z^{-n-1/2}$$

we define the normal ordering $: u(z)v(w) :$ by swapping their components:

$$(3.6) \quad : u(m)v(n) : = \begin{cases} u(m)v(n) & m < 0 \\ \frac{1}{2}(u(0)v(n) + v(n)u(0)) & m = 0 \\ v(n)u(m) & m > 0 \end{cases}$$

It then follows that $: u(z)v(w) := v(w)u(z) :$.

Based on the normal product of two fields, we can define the normal product of n fields inductively as follows. In fact we define that

$$: u_1(z_1)u_2(z_2) \cdots u_n(z_n) := : u_1(z_1)(: u_2(z_2) \cdots u_n(z_n) :) :,$$

and then use induction till we reach two fields.

We define the contraction of two states by

$$\underbrace{a(z)b(w)} = a(z)b(w) - : a(z)b(w) :,$$

which contains all poles for $a(z)b(w)$. In general, the contraction of several pairs of states is given inductively by the following rule.

THEOREM 3.1. (*Wick's theorem [FF, K]*) For elements $v_1, \dots, v_n$ we have

$$\begin{aligned} : u_1 \cdots u_m :: v_1 \cdots v_n := & : u_1 \cdots u_m v_1 \cdots v_n : + \\ & \sum_{i_1 < \cdots < i_s} \underbrace{u_{i_1} v_{j_1}} \cdots \underbrace{u_{i_s} v_{j_s}} : u_1 \cdots \hat{u}_{i_1} \cdots \hat{u}_{i_s} \cdots u_m v_1 \cdots \hat{v}_{j_1} \cdots \hat{v}_{j_s} \cdots v_n : \end{aligned}$$

where the sum runs over all possible contractions of some u_i 's and some v_i 's.

PROPOSITION 3.1. The basic operator product expansions are: for $x, y \in \mathcal{C}$ we have

$$\underbrace{u(z)v(w)} = \frac{\langle u, v \rangle}{z-w} \text{ for } \mathbb{Z} + 1/2, \quad \langle u, v \rangle \frac{(z+w)(zw)^{-1/2}}{2(z-w)} \text{ for } \mathbb{Z}.$$

In particular we have for $a, b \in P_{\mathbb{C}}$ and $\mathbb{Z} + 1/2$

$$\begin{aligned} \underbrace{a(z)b(w)} &= \underbrace{a^*(z)b^*(w)} = 0, \\ \underbrace{a(z)b^*(w)} &= \frac{\langle a, b^* \rangle}{z-w}, \quad \underbrace{a^*(z)b(w)} = \frac{\langle a^*, b \rangle}{z-w}, \end{aligned}$$

PROOF. In fact one has

$$\begin{aligned}
\underbrace{a(z)b^*(w)} &= \frac{\langle a, b^* \rangle}{2} z^{-1/2} w^{-1/2} \delta_{\mathbf{Z}, \mathbb{Z}} + \sum_{n \in \mathbf{Z}, 0 < m \in \mathbf{Z}} [a(m), b^*(n)] z^{-m-1/2} w^{-n-1/2} \\
&= \langle a, b^* \rangle \left(\frac{1}{2} z^{-1/2} w^{-1/2} \delta_{\mathbf{Z}, \mathbb{Z}} + \sum_{0 < m \in \mathbf{Z}} z^{-m-1/2} w^{m-1/2} \right) \\
&= \begin{cases} \frac{\langle a, b^* \rangle}{z-w} & \mathbf{Z} = \mathbb{Z} + 1/2 \\ \langle a, b^* \rangle \frac{(z+w)(zw)^{-1/2}}{2(z-w)} & \mathbf{Z} = \mathbb{Z} \end{cases}.
\end{aligned}$$

The other OPEs are proved in the same manner. $\square$

Note that in both cases ($\mathbb{Z}$ and $\mathbb{Z} + 1/2$) we have the following result.

PROPOSITION 3.2. *The bosonic fields satisfy the following commutation relations:*

$$[u(z), v(w)] = \langle u, v \rangle \delta(z - w)$$

PROOF. Using the fact that $: u(z)v(w) := v(w)u(z) :$ it follows from Proposition 3.1 that

$$\begin{aligned}
[u(z), v(w)] &= u(z)v(w) - v(w)u(z) \\
&= \langle u, v \rangle \delta(z - w).
\end{aligned}$$

$\square$

PROPOSITION 3.3. *The commutators among normal order products are given by:*

$$\begin{aligned}
[: a_1(z)a_2^*(z) :, : b_1(w)b_2^*(w) :] &= \langle a_1, b_2^* \rangle : a_2^*(z)b_1(z) : \delta(z - w) \\
&+ \langle a_2^*, b_1 \rangle : a_1(z)b_2^*(z) : \delta(z - w) + \langle a_1, b_2^* \rangle \langle a_2^*, b_1 \rangle \partial_w \delta(z - w).
\end{aligned}$$

PROOF. By Wick's Theorem 3.1 and Proposition 3.1, we have for $\mathbb{Z} + 1/2$

$$\begin{aligned}
: a_1(z)a_2^*(z) : : b_1(w)b_2^*(w) : &:= a_1(z)a_2^*(z)b_1(w)b_2^*(w) : \\
&+ \frac{\langle a_1, b_2^* \rangle}{z - w} : a_2^*(z)b_1(w) : + \frac{\langle a_2^*, b_1 \rangle}{z - w} : a_1(z)b_2^*(w) : + \frac{\langle a_1, b_2^* \rangle \langle a_2^*, b_1 \rangle}{(z - w)^2}.
\end{aligned}$$

Hence the result follows. The case of $\mathbb{Z}$ is shown similarly. $\square$

The (anti-symmetric) inner product of the underlying Lie algebra can be extended to that of the linear factors as follows:

$$\langle : r_1 r_2 :, : s_1 s_2 : \rangle = -\langle r_1, s_1 \rangle \langle r_2, s_2 \rangle + \langle r_1, s_2 \rangle \langle r_2, s_1 \rangle.$$

In the following we will describe the bosonic realization of the toroidal Lie algebra of classical types.

(i) Type A_{n-1} ($n \geq 2$). The roots of the finite dimensional Lie algebra are $\varepsilon_i - \varepsilon_j$ ($1 \leq i \neq j \leq n$). We associate to each finite root and imaginary root δ the

following field operators. The level $\ell = -1$.

$$\begin{aligned} X(\varepsilon_i - \varepsilon_j, z) &=: \varepsilon_i(z)\varepsilon_j^*(z) :, & h_{\varepsilon_i - \varepsilon_j}(z) &=: \varepsilon_i(z)\varepsilon_i^*(z) : - : \varepsilon_j(z)\varepsilon_j^*(z) : \\ X(\pm\alpha_i, z) &= X(\pm(\varepsilon_i - \varepsilon_{i+1}), z), & \alpha_i(z) &= h_{\varepsilon_i - \varepsilon_{i+1}}(z), \\ X(\alpha_0, z) &=: \varepsilon_n(z)\beta^*(z) :, & X(-\alpha_0, z) &=: \beta(z)\varepsilon_n^*(z) :, \\ \alpha_0(z) &=: \varepsilon_n(z)\varepsilon_n^*(z) : - : \beta(z)\beta^*(z) :. \end{aligned}$$

Using Proposition 3.3 we get for $i \neq j$

$$\begin{aligned} [X(\varepsilon_i - \varepsilon_j, z), X(\varepsilon_k - \varepsilon_l, w)] &= \delta_{jk}X(\varepsilon_i - \varepsilon_l, z)\delta(z - w) \\ &\quad - \delta_{li}X(\varepsilon_k - \varepsilon_j, z)\delta(z - w) - \delta_{jk}\delta_{li}\partial_w\delta(z - w), \\ [X(\varepsilon_i - \varepsilon_j, z), X(\varepsilon_j - \varepsilon_i, w)] &= \delta_{ij}(h_{\varepsilon_i - \varepsilon_j}(z)\delta(z - w) - \partial_w\delta(z - w)), \\ [h_{\varepsilon_i - \varepsilon_j}(z), X(\varepsilon_k - \varepsilon_l, w)] &= (\varepsilon_i - \varepsilon_j|\varepsilon_k - \varepsilon_l)X(\varepsilon_j - \varepsilon_k, w)\delta(z - w), \end{aligned}$$

where $\varepsilon_i, \varepsilon_j$ can be β (recall $(\beta|\varepsilon_i) = \delta_{1i}$). These commutation relations show that the construction provides a level -1 realization of the affine Lie algebra of type A. Moreover they also provide a representation for the toroidal Lie algebra, as the commutation relations involving α_0 are also similar to the above. In fact,

$$\begin{aligned} [X(\alpha_0, z), X(\alpha_1, w)] &=: \varepsilon_n(z)\varepsilon_2^*(z) : \delta(z - w), & [X(\alpha_0, z), X(-\alpha_1, w)] &= 0, \\ [X(\alpha_0, z), X(\alpha_n, w)] &=: - : \beta^*(z)\varepsilon_{n-1}(z) : \delta(z - w), & [X(\alpha_0, z), X(-\alpha_n, w)] &= 0. \end{aligned}$$

(ii) Type C_n ($n \geq 2$). The finite roots are $\alpha_{ij} = \frac{1}{\sqrt{2}}(\varepsilon_i - \varepsilon_j)$ ($1 \leq i \neq j \leq n$), $\pm\beta_{ij} = \pm\frac{1}{\sqrt{2}}(\varepsilon_i + \varepsilon_j)$ ($1 \leq i \neq j \leq n$). Note that $-\alpha_{ij} = \alpha_{ji}$. We realize the toroidal Lie algebra by the following field operators and the level is $\ell = -1/2$.

$$\begin{aligned} X(\alpha_{ij}, z) &=: \varepsilon_i(z)\varepsilon_j^*(z) :, & h_{\varepsilon_i - \varepsilon_j}(z) &=: \varepsilon_i(z)\varepsilon_i^*(z) : - : \varepsilon_j(z)\varepsilon_j^*(z) :, \\ X(\beta_{ij}, z) &=: \varepsilon_i(z)\varepsilon_j(z) :, & X(-\beta_{ij}, z) &=: - : \varepsilon_i^*(z)\varepsilon_j^*(z) :, \\ h_{\varepsilon_i + \varepsilon_j}(z) &=: \varepsilon_i(z)\varepsilon_i^*(z) : + : \varepsilon_j(z)\varepsilon_j^*(z) :, \\ X(\pm\alpha_i, z) &= X(\pm\alpha_{i, i+1}, z), & \alpha_i(z) &= (1/2)h_{\varepsilon_i - \varepsilon_{i+1}}(z), \\ X(\pm\alpha_n, z) &= (1/2)X(\pm\beta_{nn}, z), & \alpha_n(z) &= h_{2\varepsilon_n}(z), \\ X(\alpha_0, z) &= (1/2) : \beta^*(z)\varepsilon_1^*(z) :, & X(-\alpha_0, z) &= (1/2) : \beta(z)\varepsilon_1(z) :, \\ \alpha_0(z) &= (1/4) : \varepsilon_1(z)\varepsilon_1^*(z) : + : \beta(z)\beta^*(z) : + : \beta^*(z)\varepsilon_1(z) : + : \varepsilon_1^*(z)\beta(z) :. \end{aligned}$$

Several commutation relations are exactly the same as type A. Some new ones are obtained by Proposition 3.3 as follows.

$$\begin{aligned} [X(\beta_{ij}, z), X(-\beta_{kl}, w)] &= (\delta_{ik}X(\alpha_{jl}, z) + \delta_{il}X(\alpha_{jk}, z) + \delta_{jk}X(\alpha_{jl}, z) \\ &\quad + \delta_{jl}X(\alpha_{ik}, z))\delta(z - w) - (\delta_{ik}\delta_{jl} + \delta_{ik}\delta_{jl})\partial_w\delta(z - w), \\ [X(\alpha_{ij}, z), X(\beta_{kl}, w)] &= (\delta_{jk}X(\beta_{il}, z) + \delta_{jl}X(\beta_{ik}, z))\delta(z - w), \\ [X(\alpha_{ij}, z), X(-\beta_{kl}, w)] &= -(\delta_{ik}X(-\beta_{jl}, z) + \delta_{il}X(-\beta_{jk}, z))\delta(z - w). \end{aligned}$$

(iii) Type D_n ($n \geq 4$). The finite roots are $\pm\varepsilon_i \pm \varepsilon_j$ ($1 \leq i \neq j \leq n$). We realize the toroidal Lie algebra by the following field operators. The level $\ell = -2$.

For $i < j$ we define

$$\begin{aligned}
X(\varepsilon_i - \varepsilon_j, z) &=: \varepsilon_i(z)\varepsilon_j^*(z) : - : \varepsilon_j(z)\varepsilon_i^*(z) :, \\
h_{\varepsilon_i - \varepsilon_j}(z) &=: \varepsilon_i(z)\varepsilon_i^*(z) : - : \varepsilon_j(z)\varepsilon_j^*(z) : - : \varepsilon_i(z)\varepsilon_j^*(z) : + : \varepsilon_j(z)\varepsilon_i^*(z) :, \\
X(\varepsilon_i + \varepsilon_j, z) &=: \varepsilon_i(z)\varepsilon_j^*(z) : - : \varepsilon_j(z)\varepsilon_i^*(z) :, \\
X(-\varepsilon_i - \varepsilon_j, z) &=: \varepsilon_j(z)\varepsilon_i^*(z) : - : \varepsilon_i(z)\varepsilon_j^*(z) :, \\
h_{\varepsilon_i + \varepsilon_j}(z) &=: \varepsilon_i(z)\varepsilon_i^*(z) : + : \varepsilon_j(z)\varepsilon_j^*(z) : - : \varepsilon_i(z)\varepsilon_j^*(z) : - : \varepsilon_j(z)\varepsilon_i^*(z) :, \\
X(\pm\alpha_i, z) &= X(\pm(\varepsilon_i - \varepsilon_{i+1}), z), \quad \alpha_i(z) = h_{\varepsilon_i - \varepsilon_{i+1}}(z), \\
X(\pm\alpha_n, z) &= X(\pm(\varepsilon_{n-1} + \varepsilon_n), z), \quad \alpha_n(z) = h_{\varepsilon_{n-1} - \varepsilon_n}(z), \\
X(\alpha_0, z) &=: \varepsilon_2(z)\beta^*(z) : - : \overline{\beta}(z)\varepsilon_2^*(z) :, \\
X(-\alpha_0, z) &=: \beta^*(z)\varepsilon_2^*(z) : - : \varepsilon_2(z)\overline{\beta}(z) :, \\
\alpha_0(z) &=: \varepsilon_2(z)\varepsilon_2^*(z) : - : \varepsilon_2(z)\varepsilon_2^*(z) : + : \overline{\beta}(z)\overline{\beta}^*(z) : - : \beta^*(z)\beta(z) : .
\end{aligned}$$

(iv) Type $B_n (n \geq 2)$. The finite roots are $\pm\varepsilon_i \pm \varepsilon_j$ ($1 \leq i \neq j \leq n$) and $\pm\varepsilon_i$ ($1 \leq i \leq n$). The level $\ell = -2$. We realize the toroidal Lie algebra by the following field operators. The field operators for the roots $\pm\varepsilon_i \pm \varepsilon_j$ ($i \neq j$) are the same as in type D_n , though we need to add the bosonic fields $\varepsilon_{n+1}(z)$ and $\varepsilon_{n+1}^*(z)$ in the realization.

$$\begin{aligned}
X(\varepsilon_i, z) &=: \varepsilon_i(z)\varepsilon_{n+1}^*(z) : - : \varepsilon_{n+1}(z)\varepsilon_i^*(z) :, \\
X(-\varepsilon_i, z) &=: \varepsilon_{n+1}(z)\varepsilon_i^*(z) : - : \varepsilon_i(z)\varepsilon_{n+1}^*(z) :, \\
h_{\varepsilon_i}(z) &=: \varepsilon_i(z)\varepsilon_i^*(z) : - : \varepsilon_i(z)\varepsilon_i^*(z) :, \\
X(\pm\alpha_n, z) &= X(\pm\varepsilon_n, z), \quad \alpha_n(z) = \varepsilon_n(z),
\end{aligned}$$

and $X(\pm\alpha_i, x)$, $\alpha_i(z)$ ($0 \leq i \leq n-1$) are given by the same formulae as in type D_n . Using the same analysis we find that

$$\begin{aligned}
[X(\varepsilon_i, z), X(-\varepsilon_j, w)] &= X(\varepsilon_i - \varepsilon_j, z)\delta(z-w) - 2\delta_{ij}\partial_w\delta(z-w), \\
[X(\varepsilon_i, z), X(\varepsilon_j, w)] &= -X(\varepsilon_i + \varepsilon_j, z)\delta(z-w).
\end{aligned}$$

THEOREM 3.2. *The previous constructions give rise realizations of classical toroidal Lie algebras of types $A_{n-1} (n \geq 2)$, $C_n (n \geq 2)$, $D_n (n \geq 4)$, $B_n (n \geq 3)$ with level $-1, -1/2, -2, -2$ respectively.*

PROOF. In the above construction we have outlined major commutation relations of various types. One can use d_i to combine the argument for types A_{n-1} and C_n . We will denote $\alpha_{ij} = d_i^{-1/2}(\varepsilon_i - \varepsilon_j) = -\alpha_{ji}$ and $\beta_{ij} = d_i^{-1/2}(\varepsilon_i + \varepsilon_j)$ for $i \neq j$, then both types have the finite positive simple $\alpha_{i,i+1}$ plus $\beta_{n,n}$ in type C_n . First we note that for $i \neq j, k \neq l$

$$\begin{aligned}
[: \varepsilon_i(z)\varepsilon_j^*(z) : , : \varepsilon_k(w)\varepsilon_l^*(w) :] &= (\delta_{jk} : \varepsilon_i(z)\varepsilon_l^*(z) : \\
(3.7) \quad &- \delta_{il} : \varepsilon_j^*(z)\varepsilon_k(z) :)\delta(z-w) - \delta_{il}\delta_{jk}\partial_w\delta(z-w).
\end{aligned}$$

When $i = l, j = k$ the identity reduces to

$$[X(\alpha_{ij}, z), X(-\alpha_{ij}, w)] = d_i^{-1}(a_{ij}(z)\delta(z-w) - d_i\partial_w\delta(z-w))$$

which shows that the center $\ell = -d_i$ for type A_{n-1} and C_n . Here we also put

$$(3.8) \quad a_{ij}(z) = d_i(: \varepsilon_i(z)\varepsilon_i^*(z) : - : \varepsilon_j(z)\varepsilon_j^*(z) :).$$

Then the Heisenberg field operators are generated by the simple ones $\alpha_i(z) = a_{i,i+1}(z)$. We then have by using (3.7)

$$\begin{aligned} [\alpha_i(z), \alpha_j(w)] &= -d_i^2(2\delta_{ij} - \delta_{i+1,j} - \delta_{i,j+1})\partial_w \delta(z-w) \\ &= -d_i(\alpha_i|\alpha_j)\partial_w \delta(z-w). \end{aligned}$$

Also for any $k \neq l$ we find that (using 3.7)

$$\begin{aligned} &[\alpha_i(z), X(\alpha_{kl}, w)] \\ &= [d_i(:\varepsilon_i(z)\varepsilon_i^*(z) : - : \varepsilon_{i+1}(z)\varepsilon_{i+1}^*(z) :), : \varepsilon_k(w)\varepsilon_l^*(w)] \\ (3.9) \quad &= (\alpha_i|\alpha_{kl})X(\alpha_{kl}, z)\delta(z-w) \end{aligned}$$

Serre relations are consequences of (3.7) and (3.9) for non-special nodes. It then follows that the operators $X(\pm\alpha_i, z)$ generate a type A subalgebra for $i = 1, \dots, n-1$ ($n-2$ for A_{n-1}). The verification for special node 0 is quite similar to the above provided that we view β as ε_1 in type C_n , and β as ε_n in type A when checking commutation relations.

In types D_n and B_n , the basic commutation relations are already given above using similar analysis to Eq. (3.7). For instance in type B_n we have that

$$\begin{aligned} &[X(\alpha_0, z), X(-\alpha_1, w)] \\ &= [: \varepsilon_0(z)\varepsilon_1^*(z) : - : \varepsilon_1(z)\varepsilon_0^*(z) : , : \varepsilon_2(w)\varepsilon_1^*(w) : - : \varepsilon_1(w)\varepsilon_2^*(w) :] = 0. \end{aligned}$$

Serre relations can be easily checked from the above commutation relations plus properties of roots. We include one example to show the method. In the case of type C_n we have

$$\begin{aligned} &[X(\alpha_0, z_1), [X(\alpha_0, z_2), X(\alpha_1, w)]] = \frac{1}{2}[X(\alpha_0, z_1), [: \beta^*(z_2)\varepsilon_1^*(z_2) : , : \varepsilon_1(w)\varepsilon_2^*(w)]] \\ &= \frac{1}{4}[: \beta^*(z_2)\varepsilon_1^*(z_2) : , : \beta^*(z_2)\varepsilon_2^*(z_2) : + : \varepsilon_1^*(z_2)\varepsilon_2^*(z_2) :] \delta(z_2 - w) = 0. \\ &[X(\alpha_1, z_1), [X(\alpha_1, z_2), [X(\alpha_1, z_3), X(\alpha_0, w)]]] \\ &= \frac{1}{2}[X(\alpha_1, z_1), [X(\alpha_1, z_2), [: \varepsilon_1(z_3)\varepsilon_2^*(z_3) : , : \beta^*(w)\varepsilon_1^*(w) :]]] \\ &= -\frac{1}{2}[X(\alpha_1, z_1), [: \varepsilon_1(z)\varepsilon_2^*(z_2) : , [: \varepsilon_2^*(w)\beta^*(w) : + : \varepsilon_2^*(w)\varepsilon_1^*(w) :]]\delta(z_3 - w) \\ &= [X(\alpha_1, z_1), : \varepsilon_2^*(w)\varepsilon_2^*(w) :] \delta(z_2 - w)\delta(z_3 - w) = 0. \end{aligned}$$

□

The toroidal Lie algebra of type A_1 is also realized by the same formulae as given above. However some commutation relations are different from A_{n-1} ($n \geq 3$).

$$\begin{aligned} &[X(\alpha_0, z), X(\alpha_1, w)] = (: \varepsilon_2(w)\varepsilon_2^*(w) : - : \beta^*(w)\varepsilon_1(w) :)\delta(z-w) - \partial_w \delta(z-w), \\ &[X(-\alpha_0, z), X(-\alpha_1, w)] = (: \beta(w)\varepsilon_1^*(w) : - : \varepsilon_2(w)\varepsilon_2^*(w) :)\delta(z-w) - \partial_w \delta(z-w), \\ &[X(\alpha_0, z_1), [X(\alpha_0, z_2), X(\alpha_1, w)]] = -2 : \beta^*(w)\varepsilon_2(w) : \delta(z_1 - w)\delta(z_2 - w), \\ &[X(\alpha_1, z_1), [X(\alpha_1, z_2), X(\alpha_0, w)]] = -2X(\alpha_1, w)\delta(z_1 - w)\delta(z_2 - w), \end{aligned}$$

where the last two relations imply the Serre relations immediately for toroidal Lie algebra of type A_1 . We remark that the construction in type B_2 gives another representation of toroidal Lie algebra of type C_2 at level -2 .

References

- [BBS] S. Berman, Y. Billig, J. Szmigielski, *Vertex operator algebras and the representation theory of toroidal algebras*. Recent developments in infinite-dimensional Lie algebras and conformal field theory (Charlottesville, VA, 2000), 1–26, Contemp. Math., **297**.
- [B] Y. Billig, *Principal vertex operator representations for toroidal Lie algebras*. J. Math. Phys. **39** (1998), no. 7, 3844–3864.
- [FF] A. Feingold and I. B. Frenkel, *Classical affine algebras*, Adv. Math. **56** (1985), 117–172.
- [FJW] I. Frenkel, N. Jing, W. Wang, *Vertex representations via finite groups and the McKay correspondence* Int. Math. Res. Notices **4** (2000), 195–222.
- [G] Y. Gao, *Fermionic and bosonic representations of the extended affine Lie algebra $gl_N(\mathbb{C}_q)$* . Canad. Math. Bull. **45** (2002), no. 4, 623–633.
- [JMg] C. Jiang, D. Meng, *Vertex representations for the $\nu + 1$ -toroidal Lie algebra of type B_l* , J. Algebra **246** (2001), no. 2, 564–593.
- [JM] N. Jing, K. C. Misra, *Fermionic realization of toroidal Lie algebras of types ABD*, arXiv:0807.3056.
- [K] V. G. Kac, *Vertex algebras for beginners*, Univ. Lecture Ser., **10**, AMS, 1997.
- [L] M. Lau, *Bosonic and fermionic representations of Lie algebra central extensions*. Adv. Math. **194** (2005), no. 2, 225–245.
- [MRY] R. V. Moody, S. E. Rao, T. Yokonuma, *Lie algebras and Weyl groups arising from vertex operator representations*, Nova J. Algebra Geom. **1**(1992), no. 1, 15–57.
- [T] S. Tan, *Vertex operator representations for toroidal Lie algebra of type B_l* . Comm. Algebra **27** (1999), no. 8, 3593–3618.

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC 27695-8205, USA

E-mail address: jing@math.ncsu.edu

MISRA: DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC 27695-8205, USA

E-mail address: misra@math.ncsu.edu

XU: SCHOOL OF MATHEMATICS AND INFORMATION SCIENCE, WENZHOU UNIVERSITY, WENZHOU 325035, CHINA

E-mail address: xuchongbin1977@126.com